


Diaquatetrakis(1*H*-imidazole- $\kappa$ N<sup>3</sup>)-magnesium dichlorideM. Kayalvizhi,<sup>a</sup> G. Vasuki,<sup>a\*</sup> Kamel Kaabi<sup>b</sup> and Cherif Ben Nasr<sup>b</sup><sup>a</sup>Department of Physics, Kunthavai Naachiar Government Arts College (W) (Autonomous), Thanjavur-7, India, and <sup>b</sup>Laboratoire de Chimie des Matériaux, Faculté des Sciences de Bizerte, 7021 Zarzouna, Tunisia  
Correspondence e-mail: vasuki.arasi@yahoo.com

Received 17 July 2013; accepted 1 August 2013

Key indicators: single-crystal X-ray study;  $T = 296$  K; mean  $\sigma(\text{C}-\text{C}) = 0.002$  Å;  $R$  factor = 0.025;  $wR$  factor = 0.068; data-to-parameter ratio = 14.0.

In the title compound,  $[\text{Mg}(\text{C}_3\text{H}_3\text{N}_2)_4(\text{H}_2\text{O})_2]\text{Cl}_2$ , the  $\text{Mg}^{\text{II}}$  cation lies on a crystallographic inversion centre and is coordinated by two water molecules and four N-atom donors from monodentate imidazole ligands, giving a slightly distorted octahedral stereochemistry. In the crystal, water  $\text{O}-\text{H}\cdots\text{Cl}$  and imidazole  $\text{N}-\text{H}\cdots\text{Cl}$  hydrogen bonds give rise to a three-dimensional structure.

## Related literature

For a similar structure, see: Reiss *et al.* (2011).

## Experimental

## Crystal data

 $[\text{Mg}(\text{C}_3\text{H}_3\text{N}_2)_4(\text{H}_2\text{O})_2]\text{Cl}_2$   
 $M_r = 403.57$   
Monoclinic,  $C2/c$   
 $a = 12.3826$  (6) Å $b = 11.0048$  (4) Å  
 $c = 14.4485$  (6) Å  
 $\beta = 107.037$  (1)°  
 $V = 1882.47$  (14) Å<sup>3</sup> $Z = 4$   
Mo  $K\alpha$  radiation  
 $\mu = 0.40$  mm<sup>-1</sup> $T = 296$  K  
 $0.30 \times 0.25 \times 0.20$  mm

## Data collection

Bruker Kappa APEXII CCD diffractometer  
Absorption correction: multi-scan (SADABS; Bruker, 1999)  
 $T_{\min} = 0.889$ ,  $T_{\max} = 0.924$ 8496 measured reflections  
1854 independent reflections  
1695 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.026$ 

## Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.025$   
 $wR(F^2) = 0.068$   
 $S = 1.05$   
1854 reflections  
132 parameters  
4 restraintsH atoms treated by a mixture of independent and constrained refinement  
 $\Delta\rho_{\max} = 0.17$  e Å<sup>-3</sup>  
 $\Delta\rho_{\min} = -0.25$  e Å<sup>-3</sup>

Table 1

Selected bond lengths (Å).

Mg1—N1	2.2281 (10)	Mg1—O1	2.0923 (9)
Mg1—N3	2.1611 (10)		

Table 2

Hydrogen-bond geometry (Å, °).

$D-\text{H}\cdots A$	$D-\text{H}$	$\text{H}\cdots A$	$D\cdots A$	$D-\text{H}\cdots A$
O1—H1W $\cdots$ Cl1 <sup>i</sup>	0.84 (1)	2.30 (1)	3.1361 (9)	172 (2)
O1—H2W $\cdots$ Cl1	0.84 (1)	2.30 (1)	3.1337 (10)	176 (2)
N2—H2A $\cdots$ Cl1 <sup>ii</sup>	0.89 (1)	2.47 (1)	3.3165 (12)	160 (2)
N4—H4A $\cdots$ Cl1 <sup>iii</sup>	0.89 (1)	2.43 (1)	3.2585 (13)	155 (2)

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ ; (ii)  $x, -y+1, z-\frac{1}{2}$ ; (iii)  $-x+\frac{1}{2}, -y+\frac{1}{2}, -z+1$ .

Data collection: APEX2 (Bruker, 2004); cell refinement: APEX2 and SAINT (Bruker, 2004); data reduction: SAINT and XPREP (Bruker, 2004); program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: ORTEP-3 for Windows (Farrugia, 2012); software used to prepare material for publication: PLATON (Spek, 2009).

The authors thank the Sophisticated Analytical Instrument Facility, IIT-Madras, Chennai, for the data collection.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: ZS2271).

## References

- Bruker (1999). SADABS. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Bruker (2004). APEX2, SAINT and XPREP. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Farrugia, L. J. (2012). *J. Appl. Cryst.* **45**, 849–854.  
 Reiss, G. J., Boldog, I. & Janiak, C. (2011). *Acta Cryst.* **E67**, m1109–m1110.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Spek, A. L. (2009). *Acta Cryst.* **D65**, 148–155.

## supplementary materials

*Acta Cryst.* (2013). E69, m481 [doi:10.1107/S1600536813021478]

**Diaquatetrakis(1*H*-imidazole- $\kappa$ N<sup>3</sup>)magnesium dichloride**

**M. Kayalvizhi, G. Vasuki, Kamel Kaabi and Cherif Ben Nasr**

**1. Comment**

In the title compound,  $[\text{Mg}(\text{C}_3\text{H}_3\text{N}_2)_4(\text{H}_2\text{O})_2] \cdot 2\text{Cl}$ , the  $\text{Mg}^{\text{II}}$  cation lies on a crystallographic inversion centre and is coordinated by two water molecules and four *N*-atom donors from monodentate imidazole ligands, (Fig. 1), giving a slightly distorted octahedral geometry (Table 1). In the crystal,  $\text{O}—\text{H}\cdots\text{Cl}$  and  $\text{N}—\text{H}\cdots\text{Cl}$  hydrogen bonds between both the aqua ligands and the imidazole ligands and the chloride counter-anions (Table 2) generate a three-dimensional structure (Fig. 2). These water–chloride hydrogen-bonding interactions are in the typical range as observed in the redetermined structure of diaquatetrakis(dimethylformamide- $\kappa$ O)magnesium dichloride (Reiss *et al.*, 2011).

**2. Experimental**


A solution of  $\text{MgCl}_2$  (0.2 mmol) in water (6 ml) was added dropwise to a solution of imidazole (0.8 mmol) in ethanol. After stirring for 30 min, the mixture was filtered. Crystals suitable for X-ray analysis were obtained by evaporating the filtrate at room temperature (yield 56%).

**3. Refinement**

Carbon-bound H atoms were placed at calculated positions and treated as riding on the parent atom, with,  $\text{C}—\text{H} = 0.93 \text{ \AA}$  and with  $U_{\text{iso}}(\text{H}) = 1.2U_{\text{eq}}(\text{C})$ . The O-bound and N-bound H atoms were located in a difference Fourier map and refined freely.


**Computing details**

Data collection: *APEX2* (Bruker, 2004); cell refinement: *APEX2* and *SAINT* (Bruker, 2004); data reduction: *SAINT* and *XPREF* (Bruker, 2004); program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *ORTEP-3 for Windows* (Farrugia, 2012); software used to prepare material for publication: *PLATON* (Spek, 2009).



**Figure 1**

The molecular structure of the title compound showing atom numbering, with displacement ellipsoids drawn at the 40% probability level. For symmetry code (i):  $-x + 1/2, -y + 3/2, -z + 1$ .

**Figure 2**

Crystal packing of the title compound viewed along the *c* axis. Dashed lines indicate hydrogen bonds. H atoms not involved in hydrogen bonding have been omitted for clarity.

### Diaquatetrakis(1*H*-imidazole- $\kappa$ N<sup>3</sup>)magnesium dichloride

#### Crystal data

[Mg(C<sub>3</sub>H<sub>4</sub>N<sub>2</sub>)<sub>4</sub>(H<sub>2</sub>O)<sub>2</sub>]Cl<sub>2</sub>

*M<sub>r</sub>* = 403.57

Monoclinic, *C*2/*c*

Hall symbol: -*C* 2yc

*a* = 12.3826 (6) Å

*b* = 11.0048 (4) Å

*c* = 14.4485 (6) Å

$\beta$  = 107.037 (1)°

*V* = 1882.47 (14) Å<sup>3</sup>

*Z* = 4

*F*(000) = 840

*D<sub>x</sub>* = 1.424 Mg m<sup>-3</sup>

Mo *K*α radiation,  $\lambda$  = 0.71073 Å

Cell parameters from 8496 reflections

$\theta$  = 2.1–26.0°

$\mu$  = 0.40 mm<sup>-1</sup>

*T* = 296 K

Block, colourless

0.30 × 0.25 × 0.20 mm

#### Data collection

Bruker Kappa APEXII CCD  
diffractometer

Radiation source: fine-focus sealed tube

Graphite monochromator

$\omega$  and  $\phi$  scan

Absorption correction: multi-scan

(*SADABS*; Bruker, 1999)

*T<sub>min</sub>* = 0.889, *T<sub>max</sub>* = 0.924

8496 measured reflections

1854 independent reflections

1695 reflections with *I* > 2σ(*I*)

*R<sub>int</sub>* = 0.026

$\theta_{\max}$  = 26.0°,  $\theta_{\min}$  = 2.5°

*h* = -15→14

*k* = -13→13

*l* = -17→16

# Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.025$

$wR(F^2) = 0.068$

$S = 1.05$

1854 reflections

132 parameters

4 restraints

Primary atom site location: structure-invariant  
direct methods

Secondary atom site location: difference Fourier  
map

Hydrogen site location: inferred from  
neighbouring sites

H atoms treated by a mixture of independent  
and constrained refinement

$w = 1/[\sigma^2(F_o^2) + (0.0327P)^2 + 1.0653P]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\max} < 0.001$

$\Delta\rho_{\max} = 0.17 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\min} = -0.25 \text{ e } \text{\AA}^{-3}$

Extinction correction: *SHELXL97* (Sheldrick,  
2008),  $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$

Extinction coefficient: 0.0093 (6)

# Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

**Refinement.** Refinement of  $F^2$  against ALL reflections. The weighted  $R$ -factor  $wR$  and goodness of fit  $S$  are based on  $F^2$ , conventional  $R$ -factors  $R$  are based on  $F$ , with  $F$  set to zero for negative  $F^2$ . The threshold expression of  $F^2 > \sigma(F^2)$  is used only for calculating  $R$ -factors(gt) *etc.* and is not relevant to the choice of reflections for refinement.  $R$ -factors based on  $F^2$  are statistically about twice as large as those based on  $F$ , and  $R$ -factors based on ALL data will be even larger.

# Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
C1	0.29321 (12)	0.73665 (13)	0.29376 (10)	0.0376 (3)
H1	0.3423	0.6718	0.3154	0.045*
C2	0.19291 (13)	0.87405 (14)	0.19885 (10)	0.0440 (4)
H2	0.1594	0.9221	0.1451	0.053*
C3	0.17954 (11)	0.88354 (12)	0.28794 (9)	0.0366 (3)
H3	0.1340	0.9406	0.3059	0.044*
C4	0.01955 (12)	0.63009 (13)	0.37687 (10)	0.0393 (3)
H4	−0.0098	0.7052	0.3525	0.047*
C5	−0.03394 (12)	0.52348 (14)	0.35343 (11)	0.0462 (4)
H5	−0.1058	0.5111	0.3112	0.055*
C6	0.13086 (11)	0.49423 (12)	0.45568 (11)	0.0385 (3)
H6	0.1931	0.4549	0.4967	0.046*
N1	0.24289 (8)	0.79672 (9)	0.34828 (7)	0.0298 (2)
N2	0.26524 (11)	0.78024 (12)	0.20364 (8)	0.0434 (3)
N3	0.12407 (9)	0.61208 (9)	0.44212 (7)	0.0298 (2)
N4	0.03796 (11)	0.43755 (11)	0.40368 (10)	0.0429 (3)
O1	0.37963 (8)	0.62540 (8)	0.50646 (7)	0.0355 (2)
Mg1	0.2500	0.7500	0.5000	0.02385 (15)
Cl1	0.41012 (3)	0.35283 (3)	0.57173 (3)	0.04135 (14)
H1W	0.4370 (11)	0.6384 (16)	0.4884 (12)	0.057 (5)*
H2A	0.2880 (16)	0.7492 (16)	0.1556 (10)	0.069 (6)*
H2W	0.3849 (15)	0.5529 (10)	0.5253 (12)	0.056 (5)*
H4A	0.0289 (15)	0.3576 (9)	0.4011 (13)	0.058 (5)*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
C1	0.0447 (7)	0.0340 (7)	0.0385 (7)	0.0014 (6)	0.0188 (6)	−0.0019 (6)
C2	0.0528 (8)	0.0486 (9)	0.0284 (7)	−0.0021 (7)	0.0084 (6)	0.0050 (6)
C3	0.0413 (7)	0.0359 (7)	0.0336 (7)	0.0034 (6)	0.0125 (6)	0.0017 (6)
C4	0.0389 (7)	0.0341 (7)	0.0413 (8)	−0.0016 (6)	0.0062 (6)	0.0048 (6)
C5	0.0416 (7)	0.0472 (9)	0.0460 (8)	−0.0129 (7)	0.0071 (6)	−0.0033 (7)
C6	0.0357 (7)	0.0278 (7)	0.0535 (8)	0.0005 (5)	0.0157 (6)	0.0041 (6)
N1	0.0356 (5)	0.0277 (5)	0.0288 (5)	−0.0026 (4)	0.0138 (4)	−0.0002 (4)
N2	0.0558 (7)	0.0484 (7)	0.0325 (6)	−0.0074 (6)	0.0233 (5)	−0.0088 (5)
N3	0.0326 (5)	0.0251 (5)	0.0337 (6)	−0.0023 (4)	0.0130 (4)	0.0007 (4)
N4	0.0485 (7)	0.0261 (6)	0.0594 (8)	−0.0110 (5)	0.0244 (6)	−0.0065 (5)
O1	0.0360 (5)	0.0257 (5)	0.0523 (6)	0.0056 (4)	0.0248 (4)	0.0073 (4)
Mg1	0.0275 (3)	0.0196 (3)	0.0275 (3)	−0.0003 (2)	0.0126 (2)	0.0008 (2)
Cl1	0.0432 (2)	0.0314 (2)	0.0584 (2)	0.00885 (13)	0.02879 (17)	0.01331 (14)

*Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )*

C1—N1	1.3166 (16)	C6—N3	1.3107 (17)
C1—N2	1.3347 (18)	C6—N4	1.3306 (19)
C1—H1	0.9300	C6—H6	0.9300
C2—C3	1.3491 (19)	Mg1—N1	2.2281 (10)
C2—N2	1.355 (2)	N2—H2A	0.890 (9)
C2—H2	0.9300	Mg1—N3	2.1611 (10)
C3—N1	1.3738 (17)	N4—H4A	0.887 (9)
C3—H3	0.9300	Mg1—O1	2.0923 (9)
C4—C5	1.341 (2)	O1—H1W	0.838 (9)
C4—N3	1.3741 (17)	O1—H2W	0.839 (9)
C4—H4	0.9300	Mg1—O1 <sup>i</sup>	2.0923 (9)
C5—N4	1.355 (2)	Mg1—N3 <sup>i</sup>	2.1612 (10)
C5—H5	0.9300	Mg1—N1 <sup>i</sup>	2.2281 (10)
N1—C1—N2	111.72 (13)	C6—N3—C4	104.57 (11)
N1—C1—H1	124.1	C6—N3—Mg1	129.03 (9)
N2—C1—H1	124.1	C4—N3—Mg1	126.30 (9)
C3—C2—N2	105.92 (12)	C6—N4—C5	107.41 (12)
C3—C2—H2	127.0	C6—N4—H4A	124.5 (12)
N2—C2—H2	127.0	C5—N4—H4A	128.0 (12)
C2—C3—N1	110.17 (12)	Mg1—O1—H1W	125.7 (12)
C2—C3—H3	124.9	Mg1—O1—H2W	128.6 (12)
N1—C3—H3	124.9	H1W—O1—H2W	105.6 (17)
C5—C4—N3	110.08 (13)	O1 <sup>i</sup> —Mg1—O1	179.999 (1)
C5—C4—H4	125.0	O1 <sup>i</sup> —Mg1—N3	89.19 (4)
N3—C4—H4	125.0	O1—Mg1—N3	90.81 (4)
C4—C5—N4	106.08 (12)	O1 <sup>i</sup> —Mg1—N3 <sup>i</sup>	90.81 (4)
C4—C5—H5	127.0	O1—Mg1—N3 <sup>i</sup>	89.19 (4)
N4—C5—H5	127.0	N3—Mg1—N3 <sup>i</sup>	180.0
N3—C6—N4	111.86 (13)	O1 <sup>i</sup> —Mg1—N1 <sup>i</sup>	90.22 (4)
N3—C6—H6	124.1	O1—Mg1—N1 <sup>i</sup>	89.78 (4)

N4—C6—H6	124.1	N3—Mg1—N1 <sup>i</sup>	91.99 (4)
C1—N1—C3	104.59 (11)	N3 <sup>i</sup> —Mg1—N1 <sup>i</sup>	88.01 (4)
C1—N1—Mg1	125.70 (9)	O1 <sup>i</sup> —Mg1—N1	89.78 (4)
C3—N1—Mg1	129.45 (8)	O1—Mg1—N1	90.22 (4)
C1—N2—C2	107.60 (12)	N3—Mg1—N1	88.01 (4)
C1—N2—H2A	124.8 (12)	N3 <sup>i</sup> —Mg1—N1	91.99 (4)
C2—N2—H2A	127.5 (12)	N1 <sup>i</sup> —Mg1—N1	180.0
N2—C2—C3—N1	0.07 (16)	C4—N3—Mg1—O1 <sup>i</sup>	33.40 (11)
N3—C4—C5—N4	−0.57 (17)	C6—N3—Mg1—O1	29.08 (12)
N2—C1—N1—C3	−0.09 (15)	C4—N3—Mg1—O1	−146.60 (11)
N2—C1—N1—Mg1	174.52 (9)	C6—N3—Mg1—N1 <sup>i</sup>	−60.73 (12)
C2—C3—N1—C1	0.01 (15)	C4—N3—Mg1—N1 <sup>i</sup>	123.59 (11)
C2—C3—N1—Mg1	−174.32 (9)	C6—N3—Mg1—N1	119.27 (12)
N1—C1—N2—C2	0.13 (17)	C4—N3—Mg1—N1	−56.41 (11)
C3—C2—N2—C1	−0.12 (16)	C1—N1—Mg1—O1 <sup>i</sup>	−168.58 (11)
N4—C6—N3—C4	−0.12 (16)	C3—N1—Mg1—O1 <sup>i</sup>	4.66 (11)
N4—C6—N3—Mg1	−176.52 (9)	C1—N1—Mg1—O1	11.42 (11)
C5—C4—N3—C6	0.43 (16)	C3—N1—Mg1—O1	−175.34 (11)
C5—C4—N3—Mg1	176.96 (10)	C1—N1—Mg1—N3	−79.38 (11)
N3—C6—N4—C5	−0.23 (17)	C3—N1—Mg1—N3	93.85 (11)
C4—C5—N4—C6	0.48 (17)	C1—N1—Mg1—N3 <sup>i</sup>	100.62 (11)
C6—N3—Mg1—O1 <sup>i</sup>	−150.92 (12)	C3—N1—Mg1—N3 <sup>i</sup>	−86.14 (11)

Symmetry code: (i)  $-x+1/2, -y+3/2, -z+1$ .

### Hydrogen-bond geometry ( $\text{\AA}, ^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O1—H1 $W\cdots$ C11 <sup>ii</sup>	0.84 (1)	2.30 (1)	3.1361 (9)	172 (2)
O1—H2 $W\cdots$ C11	0.84 (1)	2.30 (1)	3.1337 (10)	176 (2)
N2—H2 $A\cdots$ C11 <sup>iii</sup>	0.89 (1)	2.47 (1)	3.3165 (12)	160 (2)
N4—H4 $A\cdots$ C11 <sup>iv</sup>	0.89 (1)	2.43 (1)	3.2585 (13)	155 (2)

Symmetry codes: (ii)  $-x+1, -y+1, -z+1$ ; (iii)  $x, -y+1, z-1/2$ ; (iv)  $-x+1/2, -y+1/2, -z+1$ .